



Referencias

Bielmann, V., Gillan, J., Perkins, N. R., Skidmore, A. L., Godden, S., y Leslie, K. E. (2010). An evaluation of Brix refractometry instruments for measurement of colostrum quality in dairy cattle. *Journal of Dairy Science*, 93(8), 3713–3721. <https://doi.org/10.3168/jds.2009-2943>

Gamsjäger, L., Elsohaby, I., Pearson, J. M., Levy, M., Pajor, E. A., Haines, D. M., y Windeyer, M. C. (2020). Assessment of Brix refractometry to estimate immunoglobulin G concentration in beef cow colostrum. *Journal of Veterinary Internal Medicine*, 34(4), 1662–1673. <https://doi.org/10.1111/jvim.15805>

Hasan, S. M. K., Junnikkala, S., Valros, A., Peltoniemi, O., y Oliviero, C. (2016). Validation of Brix refractometer to estimate colostrum immunoglobulin G content and composition in the sow. *Animal*, 10(10), 1728–1733. <https://doi.org/10.1017/S1751731116000896>

Kessler, E. C., Bruckmaier, R. M., y Gross, J. J. (2021). Short communication: Comparative estimation of colostrum quality by Brix refractometry in bovine, caprine, and ovine colostrum. *Journal of Dairy Science*, 104(2), 2438–2444. <https://doi.org/10.3168/jds.2020-19020>

Kielland, C., Rootwelt, V., Reksen, O., y Framstad, T. (2015). The association between immunoglobulin G in sow colostrum and piglet plasma. *Journal of Animal Science*, 93(9), 4453–4462. <https://doi.org/10.2527/jas.2014-8713>

Korosue, K., Murase, H., Sato, F., Ishimaru, M., Kotoyori, Y., Tsujimura, K., y Nambo, Y. (2013). Comparison of pH and refractometry index with calcium concentrations in preparturient mammary gland secretions of mares. *Journal of the American Veterinary Medical Association*, 242(2), 242–248. <https://doi.org/10.2460/javma.242.2.242>

Santiago, M. R., Fagundes, G. B., do Nascimento, D. M., Faustino, L. R., da Silva, C. M. G., Ferreira Dias, F. E., de Souza, A. P., Arrivabene, M., y Cavalcante, T. V. (2020). Use of digital Brix refractometer to estimate total protein levels in Santa Inês ewes' colostrum and lambs' blood serum. *Small Ruminant Research*, 182(September 2019), 78–80. <https://doi.org/10.1016/j.smallrumres.2019.10.014>

Silva-del-Río, N., Rolle, D., García-Muñoz, A., Rodríguez-Jiménez, S., Valldecabres, A., Lago, A., y Pandey, P. (2017). Colostrum immunoglobulin G concentration of multiparous Jersey cows at first and second milking is associated with parity, colostrum yield, and time of first milking, and can be estimated with Brix refractometry. *Journal of Dairy Science*, 100(7), 5774–5781. <https://doi.org/10.3168/jds.2016-12394>